



Health


FAMILY NAME PEARSON MRN 2276484
 GIVEN NAME OLIVER MALE FEMALE
 D.O.B. 27/03/2009 M.O. EMERGENCY DR
 ADDRESS
 LOCATION EMERGENCY DEPARTMENT

STANDARD PAEDIATRIC OBSERVATION CHART (SPOC)

5 - 11 Years

Altered Calling Criteria
 ALL OBSERVATIONS MUST BE GRAPHED

COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE



Light Blue: Increase Frequency of Observations Yellow: Clinical Review Red: Rapid Response

WP10_SI_CM03_SAGO



Health


FAMILY NAME PEARSON MRN 2276484
 GIVEN NAME OLIVER MALE FEMALE
 D.O.B. 27/03/2009 M.O. EMERGENCY DR
 ADDRESS
 LOCATION EMERGENCY DEPARTMENT

STANDARD PAEDIATRIC OBSERVATION CHART (SPOC)

5 - 11 Years

Altered Calling Criteria
 ALL OBSERVATIONS MUST BE GRAPHED

COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE



CONSIDER EARLIER ESCALATION OF PATIENTS WITH

- Chronic or complex conditions
- Post-operative
- Pre-Existing cardiac or respiratory conditions
- Opioid Infusions

ADDITIONAL CRITERIA FOR ESCALATION ON BACK PAGE

| ASSESSMENT OF RESPIRATORY DISTRESS | | | |
|------------------------------------|----------------------------------|--|--|
| | MILD | MODERATE | SEVERE |
| Airway | • Stridor on exertion | • Stridor at rest • Partial airway obstruction | • New onset of stridor • Imminent airway obstruction |
| Behaviour & Feeding | • Normal • Talks in sentences | • Some / intermittent irritability • Difficulty talking or crying • Difficulty feeding or eating | • Agitated / confused • Drowsy • Unable to talk or cry • Unable to feed or eat |
| Respiratory Rate | • Mildly increased | • Respiratory rate in the Yellow Zone | • Respiratory rate in the Red Zone • Decreasing (exhaustion) |
| Accessory Muscle Use | • None / minimal | • Moderate recession • Tracheal tug • Nasal flaring | • Severe recession • Gaspings • Grunting • Extreme pallor • Cyanosis • Absent breath sounds |
| Apnoeic Episodes | • None | • Abnormal pauses in breathing | • Apnoeic episodes |
| Oxygen | • No oxygen requirement | • Mild hypoxaemia, corrected by oxygen • Increasing oxygen requirement | • Hypoxaemia, may not be corrected by oxygen |

Holes Punched as per AS2828.1: 2012
 BINDING MARGIN - NO WRITING

SMR110018

NH606543 201213

NSW Health

STANDARD PAEDIATRIC OBSERVATION CHART (SPOC)

5 - 11 Years

Altered Calling Criteria

ALL OBSERVATIONS MUST BE GRAPHED

OTHER CHARTS IN USE

Fluid Balance Insulin Infusion Other _____
 Neurological Observation Pain / Epidural / Patient Control Analgesia Other _____
 Neurovascular Resuscitation Plan Other _____

PRESCRIBED FREQUENCY OF OBSERVATIONS

Observations must be performed routinely at least 4th hourly, unless advised below

| | | | | | |
|--------------------------------------|-------------|--|--|--|--|
| DATE: | dd/MM/yy | | | | |
| TIME: | hh:mm | | | | |
| Frequency Required | Twice daily | | | | |
| Medical Officer Name (BLOCK letters) | P. SMITH | | | | |
| Medical Officer Signature | P. SMITH | | | | |
| Attending Medical Officer Signature | R. Blagge | | | | |

ALTERATIONS TO CALLING CRITERIA
MUST BE REVIEWED WITHIN 48 HOURS OR EARLIER IF CLINICALLY INDICATED
Any alterations **MUST** be signed by a Medical Officer and confirmed by Attending Medical Officer
Document rationale for altering **CALLING CRITERIA** in the patient's health care record

| | | | | | |
|-----------------------------|-------------------|--|--|--|--|
| DATE: | dd/MM/yy | | | | |
| TIME: | hh:mm | | | | |
| Next review due Date & Time | dd/MM/yy hh:mm | | | | |

| Vital Sign | Zone | Standard Thresholds | | | | |
|------------------|-------------|----------------------|------------|--|--|--|
| Respiratory Rate | Yellow Zone | 10 - 15 35 - 50 | | | | |
| | Red Zone | <10 >50 | | | | |
| SpO ₂ | Yellow Zone | 90 - 95 | | | | |
| | Red Zone | <90 | | | | |
| Heart Rate | Yellow Zone | 60 - 70 140 - 160 | xxx-xxx | | | |
| | Red Zone | <60 >160 | ≤ or ≥ xxx | | | |
| Other | Yellow Zone | | | | | |
| | Red Zone | | | | | |

Medical Officer Name (BLOCK letters) P. SMITH

Medical Officer Signature P. SMITH

Attending Medical Officer Signature R. Blagge

| Date | Time | INTERVENTIONS / COMMENTS / ACTIONS |
|------|------|------------------------------------|
| 1. | | |
| 2. | | |
| 3. | | |
| 4. | | |

STANDARD PAEDIATRIC OBSERVATION CHART 5-11 YEARS SMR110.018

REFER TO YOUR LOCAL CLINICAL EMERGENCY RESPONSE SYSTEM (CERS) PROTOCOL FOR INSTRUCTIONS ON HOW TO MAKE A CALL TO ESCALATE CARE FOR YOUR PATIENT

CHECK THE HEALTH CARE RECORD FOR AN END OF LIFE CARE PLAN WHICH MAY ALTER THE MANAGEMENT OF YOUR PATIENT

Blue Zone Response

IF YOUR PATIENT HAS ANY BLUE ZONE OBSERVATIONS YOU MUST

1. Initiate appropriate clinical care
2. Increase the frequency of observations, as indicated by your patient's condition
3. Manage anxiety, pain and review oxygenation in consultation with the **NURSE IN CHARGE**
4. You can make a call to escalate the care of your patient at any time if you are worried or unsure whether to call

Consider the following:

1. What is usual for your patient and are there documented 'ALTERATIONS TO CALLING CRITERIA'?
2. Does the abnormal observation reflect deterioration in your patient?
3. Is there an adverse trend in observations?

Yellow Zone Response

IF YOUR PATIENT HAS ANY YELLOW ZONE OBSERVATIONS OR ADDITIONAL CRITERIA* YOU MUST

1. Initiate appropriate clinical care
2. Repeat and increase the frequency of observations, as indicated by your patient's condition
3. Consult promptly with the **NURSE IN CHARGE** to decide whether a **CLINICAL REVIEW** (or other CERS) call should be made

Consider the following:

- What is usual for your patient and are there documented 'ALTERATIONS TO CALLING CRITERIA'?
- Does the trend in observations suggest deterioration?
- Is there more than one Yellow Zone observation or additional criteria?
- Are you concerned about your patient?

IF A CLINICAL REVIEW IS CALLED:

1. Reassess your patient and escalate according to your local CERS if the call is not attended within 30 minutes or you are becoming more concerned
2. Document an A-G assessment, reason for escalation, treatment and outcome in your patient's health care record
3. Inform the Attending Medical Officer that a call was made as soon as it is practicable

***Additional YELLOW ZONE Criteria**

- Increasing oxygen requirement
- Poor peripheral circulation
- Greater than expected fluid loss
- Reduced urine output or anuria (< 1mL/kg/hr)
- Altered mental state: Agitation, Combative or Inconsolable
- New, increasing or uncontrolled pain
- New onset of fever > 38.5°C
- BGL 2-3mmol/L
- **Concern by you or any staff or family member**

CONSIDER IF YOUR PATIENT'S DETERIORATION COULD BE DUE TO SEPSIS, DEHYDRATION / HYPOVOLAEMIA / HAEMORRHAGE, OR AN OVERDOSE / OVER SEDATION

Red Zone Response

IF YOUR PATIENT HAS ANY RED ZONE OBSERVATIONS OR ADDITIONAL CRITERIA# YOU MUST CALL FOR A RAPID RESPONSE (as per local CERS) AND

1. Initiate appropriate clinical care
2. Inform the **NURSE IN CHARGE** that you have called for a Rapid Response
3. Repeat and increase the frequency of observations, as indicated by your patient's condition
4. Document an A-G assessment, reason for escalation, treatment and outcome in your patient's health care record
5. Inform the Attending Medical Officer that a call was made as soon as it is practicable

#Additional RED ZONE Criteria

- **Cardiac or respiratory arrest**
- **Circulatory collapse**
- **Patient unresponsive**
- **New onset of stridor**
- Significant bleeding
- Sudden decrease in Level of Consciousness (a drop of 2 or more points on the GCS)
- New or prolonged seizure activity
- BGL < 2mmol/L or symptomatic
- Lactate ≥ 4mmol/L
- **Serious concern by you or any staff or family member**

Holes Punched as per AS2828.1: 2012
BINDING MARGIN - NO WRITING



SMR110018